

PHOTON-DOMINATED CHEMISTRY IN THE NUCLEUS OF M82: WIDESPREAD HOC⁺ EMISSION IN THE INNER 650 PC DISK

A. FUENTE¹, S. GARCÍA-BURILLO¹, M. GERIN², D. TEYSSIER³, A. USERO^{1,4}, J.R. RIZZO^{1,5}, P. DE VICENTE¹

Not to appear in Nonlearned J., 45.

ABSTRACT

The nucleus of M82 has been mapped in several 3mm and 1mm lines of CN, HCN, C₂H, c-C₃H₂, CH₃C₂H, HC₃N and HOC⁺ using the IRAM 30m telescope. These species have been purposely selected as good tracers of photon-dominated chemistry. We have measured [CN]/[HCN]~5 in the inner 650 pc galaxy disk. Furthermore, **we have detected the HOC⁺ 1→0 line with an intensity similar to that of the H¹³CO⁺ 1→0 line. This implies a [HCO⁺]/[HOC⁺] ratio of ~ 40. These results corroborate the existence of a giant photo-dissociation region (PDR) in the nucleus of M82.** In fact, the low [HCO⁺]/[HOC⁺] ratio can only be explained if the nucleus of M82 is formed by small ($r < 0.02$ - 0.2 pc) and dense ($n \sim$ a few 10^4 - 10^5 cm⁻³) clouds immersed in an intense UV field ($G_0 \sim 10^4$ in units of the Habing field). The detection of the hydrocarbons c-C₃H₂ and CH₃C₂H in the nucleus of M82 suggests that a complex carbon chemistry is developing in this giant PDR.

Subject headings: galaxies: individual (M82) — galaxies: nuclei — galaxies: starburst — ISM: molecules — ISM: abundances — radio lines: galaxies

1. INTRODUCTION

M82 is one of the nearest and brightest starburst galaxies. Located at a distance of 3.9 Mpc, and with a luminosity of 3.7×10^{10} L_⊙, it has been extensively studied in many molecules. Compared to other prototypical nearby starburst galaxies like NGC 253 and IC 342, M82 presents systematically low abundances of the molecules NH₃, CH₃OH, CH₃CN, HNC and SiO (Takano et al. 2003). Different explanations have been proposed to account for this peculiar chemistry. Since all these molecules are related to dust grain chemistry, Takano et al. (2003) proposed that the formation of molecules on dust and/or evaporation to the gas phase is not efficient in M82.

On the other hand, several studies have revealed that the starburst has heavily influenced the interstellar medium in M82 by producing high cosmic rays and UV fluxes. A low density ionized component is filling a substantial fraction of the volume in M82 (see e.g. Seaquist et al. 1996). The molecular gas is embedded in this component in the form of warm ($T_k > 50$ K) and dense ($n > 10^4$ cm⁻³) clouds (Mao et al. 2000). Dense PDRs are expected to form in the borders of these clouds (Wolfire et al. 1990; Lord et al. 1996). Wolfire et al. (1990) have modeled the CII, SiII and OI emission and estimated a UV field of $G_0 = 10^4$ in units of the Habing field and a density of $n \sim 10^5$ cm⁻³ for the atomic component. Schilke et al. (1993) reported observations of the CI (³P₁-³P₀) line deriving a [CI]/CO column den-

sity ratio (~ 0.5) higher than that observed in non-starburst galaxies (e.g. [CI]/CO ~ 0.15 in our Galaxy). They proposed that the enhanced cosmic ray flux supplied by supernova remnants could produce the enhanced CI emission in M82. Recent results suggest that the strong UV flux dominates the chemistry of the molecular gas in M82. García-Burillo et al. (2002) obtained a high-angular-resolution image showing widespread HCO emission in this galaxy. The enhanced HCO abundance ([HCO]/[H¹³CO⁺]~3.6) measured across the whole M82 disk was interpreted in terms of a giant PDR of 650 pc size.

In this paper, we present observations of a selected set of radicals (CN, C₂H), reactive ions (CO⁺, HOC⁺) and small hydrocarbons (c-C₃H₂) which are excellent probes of the atomic to molecular transition in PDRs. In particular, the [CN]/[HCN] ratio has been successfully used as a PDR indicator in regions with very different physical conditions in our Galaxy (Fuentes et al. 1993, 1995, 1996). The detection of the reactive ion HOC⁺ is almost unambiguously associated to regions with a high ionizing flux, either PDRs or XDRs (Fuentes et al. 2003; Rizzo et al. 2003; Usero et al. 2004). Recent works have revealed that the abundances of some hydrocarbons, such as c-C₃H₂ and C₄H, are an order of magnitude larger in PDRs than those predicted by gas-phase models (Teyssier et al. 2004; Pety et al. 2004).

2. OBSERVATIONS AND ANALYSIS

The observations were carried out in June and November 2004 with the IRAM 30 m radiotelescope at Pico de Veleta (Spain). We used 2 SIS receivers tuned in single-sideband mode in the 1 mm and 3 mm bands. The observed transitions are: CN 1→0 (113.490 GHz), CN 2→1 (226.874 GHz), HCN 1→0 (88.631 GHz), C₂H 1→0 (87.317 and 87.402 GHz), c-C₃H₂ 2_{1,2}→1_{0,1} (85.339 GHz), c-C₃H₂ 6_{1,6}→5_{0,5} (217.822 GHz), CH₃C₂H 5_k→4_k (85.457 GHz), HC₃N 9→8 (81.881 GHz) and HOC⁺ 1→0 (89.487 GHz). The line intensities have been scaled

¹ Observatorio Astronómico Nacional (OAN), Apdo. 112, E-28803 Alcalá de Henares (Madrid, Spain)

² LERMA, UMR 8112, CNRS, Ecole Normale Supérieure et Observatoire de Paris, 24 rue Lhomond, 75231 Paris Cedex 05, France

³ SRON, PO Box 800, 9700 AV Groningen, The Netherlands, under an external fellowship

⁴ Instituto de Astrofísica de Andalucía (IAA), CSIC, Apdo. 3004, E-18080 Granada (Spain)

⁵ Departamento de Física, UEM, Urb. El Bosque, E-28670 Villaviciosa de Odón (Madrid, Spain)

to the main beam brightness temperature. The half-power beam width (HPBW) of the telescope is $29''$ at 85 GHz, $22''$ at 115 GHz and $12''$ at 230 GHz. We observed three positions across the M82 disk: the nucleus (RA(2000): $09^h55^m51.9^s$, Dec(2000): $69^\circ4'47.11''$) and the two peaks in the HCO emission [offsets ($+14'', +5''$) and ($-14'', -5''$) hereafter referred to as E. and W. knot respectively]. The observed spectra are shown in Fig. 1. To compare the 1.3mm and 3mm CN and c-C₃H₂ lines and make excitation calculations, we have derived beam filling factors from the H¹³CO⁺ interferometric image of García-Burillo et al. (2002). This assumption is justified since H¹³CO⁺ is an optically thin tracer of dense gas ($n > 10^4 \text{ cm}^{-3}$) and presents a quite uniform abundance in a wide range of physical conditions.

The CH₃C₂H and C₂H column densities have been estimated using the LTE approximation. For C₂H, we have assumed the typical rotation temperature in galactic PDRs, $T_{\text{rot}} = 10 \text{ K}$ (Fuente et al. 2003). In the case of CH₃C₂H, all the lines of the K-ladder are blended. In our estimates, we have assumed that 40% of the emission comes from the K=0 line and $T_{\text{rot}} = 20 \text{ K}$. These assumptions are based on the CH₃C₂H observations of Sgr B2 and Orion which have similar physical conditions to M82 (Churchwell & Hollis 1983). An LVG code has been used to derive densities and beam averaged column densities for the other species. In these calculations, we have assumed $T_k = 60 \text{ K}$ inferred from the NH₃ lines by Weiß et al. (2001).

3. RESULTS

We have estimated the hydrogen densities in the M82 disk by fitting with a LVG code the CN and C₃H₂ lines. Hydrogen densities between $n_{\text{H}_2} \sim 5 \times 10^4 \text{ cm}^{-3}$ and $n_{\text{H}_2} \sim 2 \times 10^5 \text{ cm}^{-3}$ are derived from both molecules. Assuming $n_{\text{H}_2} \sim 1 \times 10^5 \text{ cm}^{-3}$, we have calculated the column densities averaged in a beam of $29''$. The CN and HCN column densities are quite constant along the galaxy disk with values, $N(\text{CN}) = 2.0 \pm 0.5 \times 10^{14} \text{ cm}^{-2}$ and $N(\text{HCN}) = 4.0 \pm 0.5 \times 10^{13} \text{ cm}^{-2}$. The [CN]/[HCN] ratio is similar to 5 in all positions (see Table 1). As discussed in Section 4, this large value of the [CN]/[HCN] ratio is only reached in the most heavily UV exposed layers of a PDR. The derived c-C₃H₂ and CH₃C₂H column densities are also quite uniform along the disk with values, $N(\text{C}_3\text{H}_2) \sim 1.7 \pm 0.4 \times 10^{13} \text{ cm}^{-2}$ and $N(\text{CH}_3\text{C}_2\text{H}) \sim 1.0 \pm 0.6 \times 10^{14} \text{ cm}^{-2}$. These column densities are in agreement with previous estimates by Oike et al. (2004) and Mauersberger et al. (1991). Finally, we have not detected the HC₃N 9→8 line towards any position with an upper limit to the column density $N(\text{HC}_3\text{N}) < 2 \times 10^{12} \text{ cm}^{-2}$ towards the E. knot. As discussed in Section 4 the non-detection of HC₃N and the derived lower limit to the [c-C₃H₂]/[HC₃N] ratio argue in favor of a PDR chemistry in M82.

We have detected the reactive ion HOC⁺ in the three selected positions across the M82 disk. Furthermore, the intensities of the HOC⁺ 1→0 lines are similar, even larger, than those of the H¹³CO⁺ 1→0 lines. Assuming the same physical conditions for H¹³CO⁺ and HOC⁺, we derive a [HCO⁺]/[HOC⁺] ratio of ~ 40 across the 650 pc inner disk, which is two orders of magnitude lower than that found in galactic giant molecular clouds (GMCs) (Apponi & Ziurys 1997). However, the H¹³CO⁺ spec-

tra are derived from interferometric data. In order to assess the amount of missed flux (and hence the H¹³CO⁺ column density), we have compared our results to those of HCO⁺ from Nguyen-Q-Rieu et al. (1992), which were obtained by single-dish observations. From our H¹³CO⁺ spectra, we obtain $N(\text{H}^{13}\text{CO}^+) \sim 5 \times 10^{11} \text{ cm}^{-2}$ across the galaxy. Assuming a ¹²C/¹³C ratio of 89, this implies $N(\text{HCO}^+) \sim 4.5 \times 10^{13} \text{ cm}^{-2}$. This value is in agreement within a factor of 1.5 with previous estimate by Nguyen-Q-Rieu et al. (1992). Thus, even in the most conservative case we can conclude that the [HCO⁺]/[HOC⁺] ratio is < 80 in the M82 disk. Such low values of the [HCO⁺]/[HOC⁺] ratio have only been found in the galactic reflection nebula NGC 7023 (Fuente et al. 2003) and in the active nucleus of NGC 1068 (Usero et al. 2004) and put severe constraints to the ionization degree of the molecular gas in M82.

4. PHOTON-DOMINATED CHEMISTRY IN M82

Sternberg & Dalgarno (1995) found from detailed modeling that [CN]/[HCN] > 1 arises naturally in the surface layers ($A_v < 4 \text{ mag}$) of dense PDRs. Values of the [CN]/[HCN] ratio between 1–3 have been found in prototypical dense galactic PDRs (Fuente et al. 1993, 1995, 1996). We have derived [CN]/[HCN] ratios ~ 5 in all positions across the M82 nucleus suggesting that the molecular clouds in this galaxy are bathed in an intense UV field. To determine the averaged physical conditions of these clouds, we have carried out model calculations using the plane-parallel PDR model developed by Le Bourlot and collaborators (Le Bourlot et al. 1993). The model includes 135 species (HOC⁺, CH₃C₂H and HC₃N are not included) and standard gas-phase reactions. Adopting $G_0 \sim 10^4$ (Wolfire et al. 1990) and a total hydrogen nuclei density $n_{\text{H}} = n_{\text{HI}} + 2 \times n_{\text{H}_2} = 4 \times 10^5 \text{ cm}^{-3}$, the model predicts that [CN]/[HCN] ratios $\gtrsim 5$ are only expected in regions at $A_v < 5\text{--}6 \text{ mag}$ (see left panels of Fig. 2). This implies an important limit to the cloud sizes in the nucleus of M82. Since the clouds are bathed in a pervasive UV field, the averaged column density of individual clouds should be $N_{\text{H}_2} \sim 10^{22} \text{ cm}^{-2}$ in order to have averaged [CN]/[HCN] ratios ~ 5 .

Strong constraints are also derived from the HOC⁺ observations. We have measured a [HCO⁺]/[HOC⁺] ratio of ~ 40 in the M82 disk. A simple calculation of the CO⁺/HCO⁺/HOC⁺ chemical network shows that a high ionization degree, $X(\text{e}^-) > 10^{-5}$, is required to have [HCO⁺]/[HOC⁺] < 80 (Usero et al. 2004). Our PDR model shows that this high electron abundance is only found at $A_v < 4 \text{ mag}$ (see left panels of Fig. 2). This implies that the averaged column density of the clouds in the nucleus of M82 is $N(\text{H}_2) < 8 \times 10^{21} \text{ cm}^{-2}$. Thus, the M82 nucleus seems to be formed by small ($r \sim 0.02\text{--}0.2 \text{ pc}$) and dense ($n \sim 10^4\text{--}10^5 \text{ cm}^{-3}$) clouds immersed in a UV field of $G_0 \sim 10^4$. Mao et al. (2000) estimated similar column densities for the clouds in the inner 400 pc disk of M82 by modeling of the ¹²CO, ¹³CO and C¹⁸O lines. Low values of the [HCO⁺]/[HOC⁺] ratio are also found in XDRs with high ionization degree, such as active galactic nuclei (AGNs) (Usero et al. 2004). However, the PAH emission is very low in AGNs since the doubly ionized PAHs produced by X-rays are very unstable (Leach et al. 1989). The intense PAH emission observed in M82 (Normand et al. 1995; Förster Schreiber et al.

2003) indicates that X-rays are not the driving agent of the molecular gas chemistry in this galaxy.

5. HYDROCARBON CHEMISTRY

The hydrocarbon chemistry in PDRs has been a subject of increasing interest both from the theoretical and the observational point of view. Recent works have revealed that the $c\text{-C}_3\text{H}_2$ abundance in PDRs is similar to that in dark clouds. This is quite surprising taking into account that this carbon cycle is easily photodissociated. In fact, PDR gas-phase models fall short of explaining the observed $c\text{-C}_3\text{H}_2$ and C_4H abundances by an order of magnitude (Teyssier et al. 2004; Pety et al. 2004). This is clearly seen when one compares the $[c\text{-C}_3\text{H}_2]/[\text{HC}_3\text{N}]$ ratio in PDRs and dark clouds (see Table 1). While both species have similar abundances in dark clouds, the $[c\text{-C}_3\text{H}_2]/[\text{HC}_3\text{N}]$ ratio is >10 in PDRs. Since both molecules are easily destroyed by photodissociation, this suggests the existence of an additional $c\text{-C}_3\text{H}_2$ formation mechanism in PDRs. Several authors have proposed formation processes of small hydrocarbons linked to the PAHs chemistry (Teyssier et al. 2004; Pety et al. 2004). We have detected widespread emission of C_2H and $c\text{-C}_3\text{H}_2$ in the M82 disk. However, we have not detected HC_3N towards any position. The derived upper limits to the HC_3N column density in M82 show that the $[c\text{-C}_3\text{H}_2]/[\text{HC}_3\text{N}]$ ratio is at least a factor of 10 larger in this starburst galaxy than in the prototypical galactic dark cloud TMC 1, and is similar to that found in dense galactic PDRs (see Table 1). This result reinforces the scenario where the inner 650 pc region of M82 is a giant PDR.

In addition to C_2H and $c\text{-C}_3\text{H}_2$, we have detected $\text{CH}_3\text{C}_2\text{H}$ in all the positions across the M82 disk. This detection is rather puzzling since this molecule is easily photodissociated in regions exposed to intense UV fields and therefore its abundance is expected to be very low in these regions. Furthermore, this molecule is not detected in prototypical galactic PDRs, like the Horsehead, which is known to be specially rich in carbon compounds (Teyssier et al. 2004; Pety et al. 2004). One possibility is that the $\text{CH}_3\text{C}_2\text{H}$ emission arises in a population of clouds similar to the galactic GMCs. The molecule $\text{CH}_3\text{C}_2\text{H}$ is quite abundant in GMCs, presenting similar abundances to those of chemically related species like CH_3OH and CH_3CN . The same behavior is observed in other prototypical starburst galaxies like NGC 253 where $[\text{CH}_3\text{C}_2\text{H}]/[\text{CH}_3\text{OH}] \sim 1$ and $[\text{CH}_3\text{C}_2\text{H}]/[\text{CH}_3\text{CN}] \sim 8$. However, $[\text{CH}_3\text{C}_2\text{H}]/[\text{CH}_3\text{OH}] > 8$ and $[\text{CH}_3\text{C}_2\text{H}]/[\text{CH}_3\text{CN}] > 25$ in M82 (Mauersberger et al. 1991; Huettemeister et al. 1997), i.e., a factor 3–10 larger than in NGC 253. These abundance ratios show that the hydrocarbon chemistry in M82 is very different from that of GMCs.

We think that the over-abundance of $\text{CH}_3\text{C}_2\text{H}$ is also related to the enhanced UV flux in this galaxy. In fact, $\text{CH}_3\text{C}_2\text{H}$ and $\text{CH}_3\text{C}_4\text{H}$ has been detected by Cernicharo et al. (2001) in the C-rich proto-planetary nebula CRL 618. This nebula is specially rich in complex carbon compounds, in particular, polyynes ($\text{C}_{2n+2}\text{H}_2$) and methylpolyynes ($\text{CH}_3\text{C}_{2n}\text{H}$). They argued that the formation of these complex molecules has started by the photolysis of acetylene (C_2H_2) and methane (CH_4), both processes occurring in the inner PDR of the planetary

nebula. The photo-erosion of PAHs could also contribute to enhance the abundance of these molecules by releasing small carbon chains and acetylene to the gas phase.

6. A GLOBAL SCENARIO FOR THE CHEMISTRY IN M82

The observations presented in this paper show that the strong UV flux drives the chemistry of the molecular gas in the M82 disk. In this Letter, we explore whether the PDR chemistry can solely account for all the molecular abundances measured in this galaxy so far. For this aim, we compare the molecular abundances in M82 to those of the prototypical starburst NGC 253. Molecules such as NH_3 , CH_3OH , CH_3CN , SO , SiO and HC_3N are clearly under-abundant in M82 compared to NGC 253 (Weiß et al. 2001; Huettemeister et al. 1997; Mauersberger et al. 1991; Takano, et al. 1995; García-Burillo et al. 2000, 2001). All these molecules are easily photodissociated and are not expected in a PDR. Other species such as CS , CN , HCN and HCO^+ present fractional abundances $\sim$ a few 10^{-9} in M82 which are very similar to those measured in NGC 253 (Mauersberger & Henkel 1989). These molecules belong to the reduced group of species which have significant abundances at $A_v < 5$ mag in our PDR model (see left panels of Fig. 2). However, our model predicts abundances a factor of 10 lower than those observed. Cosmic rays and/or shocks could contribute to enhance the abundances of these species.

To investigate the possible effect of the enhanced cosmic ray flux on the chemistry, we have repeated the model calculations with an enhanced cosmic rays flux of $\zeta = 4 \times 10^{-15} \text{ s}^{-1}$. This is the value estimated by Suchkov et al. (1993) to account for the physical conditions of the molecular gas in M82 and is, very likely, an upper limit to the actual cosmic ray flux. The enhanced cosmic rays flux does not change significantly the value of the $[\text{CN}]/[\text{HCN}]$ ratio and the ionization degree at $A_v < 5$ mag (see right panels of Fig. 2). However, the fractional abundances of HCN , CN and CS averaged over the region at $A_v < 5$ mag are a factor of 10 larger than those obtained with the standard value $\zeta = 5 \times 10^{-17} \text{ s}^{-1}$. Moreover, these values are in reasonable agreement with the observational results in M82. Thus, by including the enhanced cosmic ray flux we get a better fit of the model to the observational results. However, the HCO^+ abundance still remains a factor of 10 lower than the observed one.

Some of the underabundant species in M82 like SiO are well known shock chemistry tracers. This suggests that shocks are not the dominant mechanism at work in the galaxy disk. The SiO map reported by García-Burillo et al. (2001) show that shocks are not occurring in the galaxy disk but in the disk-halo interface because of the hot gas coming out of the plane in the nucleus of M82. This is consistent with the interpretation of M82 as an evolved starburst compared to NGC 253. While in NGC 253 the molecular chemistry is determined by the shocks associated with the early stages of star formation, in M82 it is determined by the UV radiation produced by evolved stars and the enhanced cosmic rays flux (García-Burillo et al. 2002).

We are grateful to the technical staff in Pico de Veleta for their supporting during observations. This paper

TABLE 1
RELATIVE FRACTIONAL ABUNDANCES

Molecule	M82		Orion Bar		NGC 7023	Horsehead	TMC1	L134N
	E.	(0,0)	W.	IF	PDR peak	IR peak	CP	
C ₂ H (10 ¹³ cm ⁻³)	58	41	30	57 ^(a)	3.8 ^(d)	16 ^(e)	7.2 ^(f)	15 ^(g)
c-C ₃ H ₂ /C ₂ H	0.02	0.04	0.02	0.02 ^(b)	0.03 ^{(b)(d)}	0.06 ^(e)	0.8 ^(g)	0.33 ^(g)
CH ₃ C ₂ H/C ₂ H	0.2	0.4	0.3	...	...	<0.07 ^(e)	1.4 ^(f)	0.07 ^(g)
c-C ₃ H ₂ /HC ₃ N	> 6	> 4	> 3	>10 ^{(b)(c)}	...	19 ^(e)	0.4 ^(h)	3 ^{(g),(i)}
CN/HCN	4	6	4	3 ^(a)	4-8 ^(d)	...	0.07 ^(f)	0.07 ⁽ⁱ⁾
CN/HC ₃ N	>120	>58	>80	> 180 ^(a)	...	...	0.05 ^{(f)(h)}	0.9 ⁽ⁱ⁾
HCO ⁺ /HOC ⁺	35	49	50	<166 ^(a)	50-120 ^(b)	...	...	...

REFERENCES. — (a) Fuente et al.(1996); (b) Fuente et al.(2003); (c) Rodríguez-Franco et al.(1998); (d) Fuente et al.(1993); (e) Teyssier et al.(2004); (f) Patrap et al.(1997); (g) Fossé (2003); (h) Takano et al. 1998; (i) Dickens et al. (2000).

has been partially funded by the Spanish MCyT under projects DGES/AYA2000-927, ESP2001-4519-PE, ESP2002-01693, AYA2002-01241 and AYA2003-06473.

We are also grateful to Jacques Le Bourlot for his important help with model calculations.

REFERENCES

- Apponi, A. J. & Ziurys, L. M. 1997, ApJ, 481, 800
Cernicharo, J., Heras, A. M., Tielens, A. G. G. M., Pardo, J. R., Herpin, F., Guélin, M., & Waters, L. B. F. M. 2001, ApJ, 546, L123
Churchwell, E. & Hollis J.M., 1983, ApJ, 272, 591
Dickens, J. E., Irvine, W. M., Snell, R. L., Bergin, E. A., Schloerb, F. P., Pratap, P., & Miralles, M. P. 2000, ApJ, 542, 870
Fossé, D., 2003, Ph. D. dissertation, Université Pierre et Marie Curie, Paris, France
Förster Schreiber N.M., Sauvage M., Charmandaris V. et al., 2003, A&A, 399, 833
Fuente, A., Martín-Pintado, J., Cernicharo, J., & Bachiller, R. 1993, A&A, 276, 473
Fuente, A., Martín-Pintado, J., & Gaume, R. 1995, ApJ, 442, L33
Fuente, A., Rodríguez-Franco, A., & Martín-Pintado, J. 1996, A&A, 312, 599
Fuente, A., Rodríguez-Franco, A., García-Burillo, S., Martín-Pintado, J., & Black, J. H. 2003, A&A, 406, 899
García-Burillo, S., Martín-Pintado, J., Fuente, A., & Neri, R. 2000, A&A, 355, 499
García-Burillo, S., Martín-Pintado, J., Fuente, A., & Neri, R. 2001, ApJ, 563, L27
García-Burillo, S., Martín-Pintado, J., Fuente, A., Usero, A., & Neri, R. 2002, ApJ, 575, L55
Huettemeister, S., Mauersberger, R., & Henkel, C. 1997, A&A, 326, 59
Leach, S., Eland, J.H.D., Price, S.D., 1989, J. Phys. Chem. 93, 7583
Le Bourlot, J., Pineau Des Forets, G., Roueff, E., & Flower, D. R. 1993, A&A, 267, 233
Lord S.D., Hollenbach D.J., Haas M.R., Rubin R.H., Colgan S.W.J., Erickson E.F. 1996, ApJ, 465, 703
Mao R.Q., Henkel C., Schulz A. et al. 2000, A&A 358, 433
Mauersberger, R. & Henkel, C. 1989, A&A, 223, 79
Mauersberger, R., Henkel, C., Walmsley, C. M., Sage, L. J., & Wiklund, T. 1991, A&A, 247, 307
Normand P., Rouan D., Lacombe F., Tiphène D. 1995, A&A, 297, 311
Nguyen, Q.-R., Jackson, J. M., Henkel, C., Truong, B., & Mauersberger, R. 1992, ApJ, 399, 521
Oike, T., Kawaguchi, K., Takano, S., & Nakai, N. 2004, PASJ, 56, 431
Pety J., Teyssier D., Fossé D., Gerin M. et al., 2004, submitted to A&A
Pratap, P., Dickens, J. E., Snell, R. L., Miralles, M. P., Bergin, E. A., Irvine, W. M., & Schloerb, F. P. 1997, ApJ, 486, 862
Rizzo, J. R., Fuente, A., Rodríguez-Franco, A., & García-Burillo, S. 2003, ApJ, 597, L153
Rodríguez-Franco, A., Martín-Pintado, J., & Fuente, A. 1998, A&A, 329, 1097
Schilke, P., Carlstrom, J. E., Keene, J., & Phillips, T. G. 1993, ApJ, 417, L67
Sequist, E. R., Carlstrom, J. E., Bryant, P. M., & Bell, M. B. 1996, ApJ, 465, 691
Suchkov, A., Allen, R. J., & Heckman, T. M. 1993, ApJ, 413, 542
Sternberg, A. & Dalgarno, A. 1995, ApJS, 99, 565
Takano, S., Nakai, N., & Kawaguchi, K. 1995, PASJ, 47, 801
Takano, S., et al. 1998, A&A, 329, 1156
Takano, S., Nakai, N., Kawaguchi, K., Takano, T., Schilke, P., & Winnewisser, G. 2003, The Astrochemistry of External Galaxies, 25th meeting of the IAU, Joint Discussion 21, 23 July 2003, Sydney, Australia, 21
Teyssier, D., Fossé, D., Gerin, M., Pety, J., Abergel, A., & Roueff, E. 2004, A&A, 417
Usero, A., García-Burillo, S., Fuente, A., Martín-Pintado, J., & Rodríguez-Fernández, N. J. 2004, A&A, 419, 897
Weiß, A., Neininger, N., Henkel, C., Stutzki, J., & Klein, U. 2001, ApJL, 554, L143
Wolfire, M. G., Tielens, A. G. G. M., & Hollenbach, D. 1990, ApJ, 358, 116




FIG. 1.— Observed spectra towards the positions East (E), Center and West (W) in M82. The 30m beam at 90 GHz around the three observed positions has been drawn in the interferometric HCO image by García-Burillo et al. (2002). We have fully sampled the HCO disk at 3mm.




FIG. 2.— Predictions for the abundances of various species derived from updated Le Bourlot et al.'s model. The calculations have been carried out for $n=4 \times 10^5 \text{ cm}^{-3}$ and $G_0=1 \times 10^4$ in units of the Habing field. The cosmic ray flux is set to $\zeta=5 \times 10^{-17} \text{ s}^{-1}$ (galactic value) in left panels and $\zeta=4 \times 10^{-15} \text{ s}^{-1}$ (M82 value as derived by Suchkov et al. 1993) in right panels. We have shadowed the region of the plot in agreement with the observational results in M82.